

## Scientific Paper:

The Journal of Experimental Biology, 207, 3629-3637, 2004

# The effects of sustained exercise and hypoxia upon oxygen tensions in the red muscle of rainbow trout

D. J. McKenzie<sup>1,\*</sup>, S. Wong<sup>2</sup>, D. J. Randall<sup>3</sup>, S. Egginton<sup>4</sup>, E. W. Taylor<sup>1</sup> and A. P. Farrell<sup>2</sup>

<sup>1</sup>School of Biosciences, University of Birmingham, Birmingham B15 2TT, UK,

<sup>2</sup>Department of Biological Sciences, Simon Fraser University, 8888 University Drive, Burnaby, BC, V5A 1S6, Canada

<sup>3</sup>Department of Biology and Chemistry, City University of Hong Kong, Tat Chee Avenue, Kowloon, Hong Kong, China

<sup>4</sup>Department of Physiology, University of Birmingham, Birmingham B15 2TT, UK

\*Author for correspondence at present address: CNRS/IFREMER, CREMA L'Houmeau, BP 5, 17137 L'Houmeau, France [e-mail: David.Mckenzie@ifremer.fr]

### Abstract:

Teleost fish possess discrete blocks of oxidative red muscle (RM) and glycolytic white muscle, whereas tetrapod skeletal muscles are mixed oxidative/glycolytic. It has been suggested that the anatomy of RM in teleost fish could lead to higher intramuscular O<sub>2</sub> partial pressures ( $P_{O_2}$ ) than in mammalian skeletal muscles. This study provides the first direct experimental support for this suggestion by using novel optical fibre sensors to discover a mean ( $\pm$  S.E.M.,  $N=6$ ) normoxic steady-state red muscle  $P_{RM_{O_2}}$  of  $61\pm 10$  mmHg (1 mmHg=133.3 Pa) in freeswimming rainbow trout *Oncorhynchus mykiss*. This is significantly higher than literature reports for mammalian muscles, where the  $P_{O_2}$  never exceeds 40 mmHg. Aerobic RM powers sustained swimming in rainbow trout. During graded incremental exercise,  $P_{RM_{O_2}}$  declined from  $62\pm 5$  mmHg at the lowest swim speed down to  $45\pm 3$  mmHg at maximum rates of aerobic work, but then rose again to  $51\pm 5$  mmHg at exhaustion. These measurements of  $P_{RM_{O_2}}$  during exercise indicated, therefore, that O<sub>2</sub> supply to the RM was not a major limiting factor at exhaustion in trout. The current study found no evidence that teleost haemoglobins with a Root effect cause extremely elevated O<sub>2</sub> tensions in aerobic tissues. Under normoxic conditions,  $P_{RM_{O_2}}$  was significantly lower than arterial  $P_{O_2}$  ( $119\pm 5$  mmHg), and remained lower when the arterial to tissue  $P_{O_2}$  gradient was reduced by exposure to mild hypoxia. When two sequential levels of mild hypoxia (30 min at a water  $P_{O_2}$  of 100 mmHg then 30 min at 75 mmHg) caused  $P_{a_{O_2}}$  to fall to  $84\pm 2$  mmHg then  $61\pm 3$  mmHg, respectively, this elicited simultaneous reductions in  $P_{RM_{O_2}}$ , to  $51\pm 6$  mmHg then  $41\pm 5$  mmHg, respectively. Although these hypoxic reductions in  $P_{RM_{O_2}}$  were significantly smaller than those in  $P_{a_{O_2}}$ , the effect could be attributed to the sigmoid shape of the trout haemoglobin–O<sub>2</sub> dissociation curve.

Key-words: O<sub>2</sub>-sensitive optode, Root effect, O<sub>2</sub> partial pressure, arterial blood O<sub>2</sub> content, O<sub>2</sub> consumption, swimming