

Scientific Paper:

Wiley Periodicals, Inc. DOI: 10.1002, bit.20812, 2006

Competition Between Oxygen and Nitrate Respirations in Continuous Culture of *Pseudomonas aeruginosa* Performing Aerobic Denitrification

Fan Chen, Qing Xia, Lu-Kwang Ju

Department of Chemical Engineering, The University of Akron, Akron, Ohio 44325-3906; telephone: 330-972-7252; fax: 330-972-5856; e-mail: LukeJu@Uakron.edu

Abstract:

Continuous culture of *P. aeruginosa* was conducted with nitrate-containing media under the dilution rates (D) of 0.026, 0.06, and 0.13/h and the dissolved oxygen concentrations (DO) of 0–2.2 mg/L. The bacterium performed simultaneous O_2 and nitrate respiration in all of the systems studied. For each D , the (apparent) cell yield from glucose ($Y_{X/S}$) was lower at zero DO , but did not change substantially with non-zero DO . In non-zero DO systems, $Y_{X/S}$ increased with increasing D , and when fit with a model considering cell death, gave the following parameters: maximum cell yield $Y_{X/S}^m \approx 0.49$, maintenance coefficient $M_s = 0.029$ (/h), and cell decay constant $k_d = 0.014$ /h. The same model failed to describe the behaviors of zero- DO systems, where neither glucose nor nitrate was limiting and the limiting factor(s) remained unknown. The cell yield from accepted electron ($Y_{X/e}$) was however relatively constant in all systems, and the energy yield per electron accepted via denitrification was estimated at $\sim 69\%$ of that via O_2 respiration. A closer examination revealed that increasing DO enhanced O_2 respiration only at extremely low DO (< 0.05 mg/L), beyond which the increasing DO only slightly increased its weak inhibition on denitrification. While O_2 was the preferred electron acceptor, the fraction of electrons accepted via denitrification increased with increasing D .

Key-words: *Pseudomonas aeruginosa*, respiration, denitrification, cell yield, energy yield