

Scientific Paper:

Marine Ecology Progress Series 366, 305-309, 2008

Oxygen consumption of a single embryo/planula in the reef-building coral *Acropora intermedia*

N. Okubo^{1,5,*}, H. H. Yamamoto², F. Nakaya³, K. Okaji⁴

¹Graduate School of Environment and Information Sciences, Yokohama National University, Tokiwadai 79-2, Hodogaya, Yokohama 240-8501, Japan

²Okinawa Churaumi Aquarium, Motobu-cho, Okinawa 905-0206, Japan

³Graduate School of Humanities and Science, Ochanomizu University, Bunkyo-ku, Tokyo 112-8610, Japan

⁴CoralQuest Inc., Asahicho 1-34-10, Atsugi, Kanagawa 243-0014, Japan

⁵Present address: Japan Society for the Promotion of Science/Seto Marine Biological Laboratory, Field Science Education and Research Center, Kyoto University, Shirahama, Nishimuro, Wakayama 649-2211, Japan

Abstract:

O₂ consumption of a single embryo/planula at each developmental stage was monitored in the reef-building coral *Acropora intermedia* using an optical O₂-sensing system with our original micro-chamber system (6.28 μl). The lowest rate of O₂ consumption was observed in unfertilized eggs. After fertilization, O₂ consumption increased and remained constant until the prawn chip blastula stage. However, O₂ consumption began to increase again during the bowl-shaped blastula stage, which involves the formation of 2 germ layers and corresponds to the beginning of gastrulation. The rate of O₂ consumption peaked during the teardrop-shaped planula stage. During this stage planulae are able to swim actively, especially in the vertical plane, so an increase in energy consumption during this stage is to be expected. O₂ consumption began to decrease gradually 5 d after spawning. At this stage, the larvae frequently touched the substrate with their concave aboral end, which features numerous spirocysts required for substrate attachment. When the planulae began to settle, 7 d after spawning, the rate of O₂ consumption dropped to that of unfertilized eggs, suggesting that the planulae slowly use stored energy for crawling/settlement behavior and/or post-settlement growth and survivorship.

Key-words: Development, dispersal, energy, larva, lecithotrophic, metabolism, recruitment, settlement, competency period