

Scientific Paper:

J Comp Physiol B 177, 579-587, 2007

Continuous measurement of oxygen tensions in the air-breathing organ of Pacific tarpon (*Megalops cyprinoides*) in relation to aquatic hypoxia and exercise

Roger S. Seymour¹, Anthony P. Farrell², Keith Christian³, Timothy D. Clark¹, Michael B. Bennett⁴, Rufus M. G. Wells⁵, John Baldwin⁶

¹Environmental Biology, School of Earth and Environmental Sciences, University of Adelaide, Adelaide 5005, Australia; e-mail: roger.seymour@adelaide.edu.au

²Department of Zoology and Faculty of Land and Food Systems, University of British Columbia, Vancouver, BC, Canada V6T 1Z4

³School of Science, Charles Darwin University, Darwin, NT 0909, Australia

⁴School of Biomedical Sciences, The University of Queensland, St. Lucia, QLD 4072, Australia

⁵School of Biological Sciences, The University of Auckland, Private Bag 92019, Auckland, New Zealand

⁶School of Biological Sciences, Monash University, Clayton Campus, VIC 3800, Australia

Abstract:

The Pacific tarpon is an elopomorph teleost fish with an air-breathing organ (ABO) derived from a physostomous gas bladder. Oxygen partial pressure (PO_2) in the ABO was measured on juveniles (238 g) with fiberoptic sensors during exposure to selected aquatic PO_2 and swimming speeds. At slow speed (0.65 BL s^{-1}), progressive aquatic hypoxia triggered the first breath at a mean PO_2 of 8.3 kPa. Below this, opercular movements declined sharply and visibly ceased in most fish below 6 kPa. At aquatic PO_2 of 6.1 kPa and swimming slowly, mean airbreathing frequency was 0.73 min^{-1} , ABO PO_2 was 10.9 kPa, breath volume was 23.8 ml kg^{-1} , rate of oxygen uptake from the ABO was $1.19 \text{ ml kg}^{-1} \text{ min}^{-1}$, and oxygen uptake per breath was 2.32 ml kg^{-1} . At the fastest experimental speed (2.4 BL s^{-1}) at 6.1 kPa, ABO oxygen uptake increased to about $1.90 \text{ ml kg}^{-1} \text{ min}^{-1}$, through a variable combination of breathing frequency and oxygen uptake per breath. In normoxic water, tarpon rarely breathed air and apparently closed down ABO perfusion, indicated by a drop in ABO oxygen uptake rate to about 1% of that in hypoxic water. This occurred at a wide range of ABO PO_2 (1.7–26.4 kPa), suggesting that oxygen level in the ABO was not regulated by intrinsic receptors.

Key-words: Fish; Respiration; Air-breathing; Bimodal gas exchange; Oxygen receptors